

BALL STATE UNIVERSITY Executive Summary Fall 2024

American College Health Association National College Health Assessment III

ACHA-NCHA III

The ACHA-NCHA III supports the health of the campus community by fulfilling the academic mission, supporting short- and long-term healthy behaviors, and gaining a current profile of health trends within the campus community.



Copyright 2024 All Rights Reserved

Table of Contents

II. Findings	
A. General Health and Campus Climate	
B. Nutrition, BMI, Physical Activity, and Food Security	
C. Healthcare Utilization	
D. Impediments to Academic Performance	
E. Violence, Abusive Relationships, and Personal Safety	
F. Tobacco, Alcohol, and Other Drug Use	
G. Sexual Behavior	
H. Mental Health and Wellbeing	
I. Acute Conditions	
J. Ongoing or Chronic Conditions	
K. Sleep	

ACHA, the nation's principal advocate and leadership organization for college and university health, represents a diverse membership that provides and supports the delivery of healthcare and prevention and wellness services for the nation's 20 million college students. For more information about the association's programs and services, visit www.acha.org, and www.acha.org/NCHA.

Suggested citation for this document:

American College Health Association. American College Health Association-National College Health Assessment III: Ball State University Executive Summary Fall 2024. Silver Spring, MD: American College Health Association; 2024.

Introduction and Notes

The ACHA-National College Health Assessment (ACHA-NCHA) is a national research survey organized by the American College Health Association (ACHA) to assist college health service providers, health educators, counselors, and administrators in collecting data about their students' habits and behaviors on the most prevalent health topics. The ACHA-NCHA now provides the largest known comprehensive data set on the health of college students, providing the college health and higher education fields with a vast spectrum of information on student health.

ACHA initiated the original ACHA-NCHA in 2000 and the instrument was used nationwide through the Spring 2008 data collection period. A revised survey, the ACHA-NCHA-II, was in use from Fall 2008 - Spring 2019 data collection periods. The survey was redesigned again, and data collection with the ACHA-NCHA III began in Fall 2019.

Please note that it is not appropriate to compare trends between versions of the survey. Directly comparing data points between the Original ACHA-NCHA, the ACHA-NCHA II, and the ACHA-NCHA III can lead to an erroneous conclusion and is not recommended.

Notes about this report:

- 1. Missing values have been excluded from analysis and only valid percents are included in this document, unless otherwise noted.
- 2. The ACHA-NCHA III is programmed differently than earlier versions of the survey. Rather than asking the respondents to answer every question (and offering a "not applicable" option), display logic was used throughout the survey to determine whether, based on their response to an earlier question, the student saw a follow-up question. This makes the valid percents of certain questions impossible to apply to the entire sample, as the denominator used was limited to only the number of students that saw the question. When appropriate, results are also presented using the entire sample as the denominator to show the proportion of the overall sample that experienced a particular issue. These differences in presentation are carefully noted throughout the document and will often explain differences observed between this document and the full data report. Please look carefully at descriptions of the data presented in each table, as well as any footnotes included.
- 3. About the use of sex and gender in this report: Survey results are reported by sex based on the responses to questions 67A, 67B, and 67C. The responses to these questions are used to create a new variable called RSEX. RSEX is used for organizing results in the ACHA-NCHA report documents. Respondents are reported as cis men or cis women only when their responses to 67A, 67B, and 67C are consistent with one another. If gender identity is consistent with sex at birth AND "no" is selected for transgender, then respondents are designated as either cis men or cis women in RSEX. If respondents select "yes" for transgender OR their sex at birth is not consistent with their gender identity, then they are designated as transgender/gender non-conforming in RSEX. A respondent that selects "intersex" for sex at birth, "no" for transgender, and man or woman for gender identity are designated as cis men or cis women in RSEX. A respondent that selects "intersex" for sex at birth, "yes" for transgender, or selects a gender identity other than man or woman are designated as transgender/gender non-conforming in RSEX. A respondent that selects "another identity" on 67C is designated missing in RSEX. A respondent that skips any of the three questions is designated as missing in RSEX. Totals displayed in this report include missing responses. Please see the ACHA-NCHA III survey codebook for more information about how data on sex and gender are coded.

Beginning in Spring 2021, responses for transgender and gender-nonconforming students became readily available directly in the report documents. We've prepared the following tips for using these report documents.

Important considerations with this format:

- Percentages in the Executive Summary may represent a very small number of transgender/gender non-conforming students and can limit the generalizability of a particular finding. To assist with the interpretation of the percentages displayed in the Executive Summary, the total sample size for each group has been added to every page.
- We encourage ACHA-NCHA surveyors to carefully review their report documents, particularly among the student demographic variables, and consider students who may be inadvertently identified in the results based on a unique combination of the demographic characteristics before sharing the documents widely or publicly. This is especially true for very small schools, as well as schools that lack diversity in the student population.
- Think about the implication of working with and documenting very small samples from the perspective of making meaningful interpretations, as well as the privacy of respondents. This is true of all demographic variables, and not limited to gender identity. You may consider a minimum cell size or another threshold by which you make decisions about making your Institutional Data Report publicly available. It is less of a concern in your Institutional Executive Summary as we only display the percentages with the overall sample size.
- 4. To see a list of changes made to the current and past surveys, click here: https://www.acha.org/ncha/resources/resources-for-participating-institutions/current-survey/.

For additional information about the survey's development, design, and methodology, email Mary T Hoban, PhD, MCHES, (mhoban@acha.org), Christine Kukich, MS (ckukich@acha.org), or visit www.acha-ncha.org.

This Executive Summary highlights results of the ACHA-NCHA III Fall 2024 survey for Ball State University consisting of 1117 respondents. The response rate was 9%.

Findings

A. General Health and Campus Climate

Cis Men n = 239
Cis Women n = 742
Trans/GNC n = 124

• 38.1 % of college students surveyed (50.2 % cis men, 37.8 % cis women, and 19.3 % transgender/gender non-conforming) described their health as *very good or excellent*.

• 81.5 % of college students surveyed (86.2 % cis men, 82.9 % cis women, and 64.9 % transgender/gender non-conforming) described their health as *good*, *very good*, *or excellent*.

Proportion of college students who reported they agree or strongly agree that:			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
Percent (%)		conforming	
I feel that I belong at my college/university	73.1	67.5	62.1	68.0
I feel that students' health and well-being are a priority of my college/university	48.7	46.1	33.1	44.8
I feel that the climate of my college/university encourages free and open discussion about students				
health and well-being	63.7	59.2	51.6	59.1
We are a college/university where we look out for each other	46.0	46.0	38.7	45.0

B. Nutrition, BMI, Physical Activity, and Food Security

				Trans/	
College students reported:		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Drinking 0 sugar-sweetened beverages (per day), on average, in the last 7 days		24.7	15.6	17.1	18.1
Drinking 1 or more sugar-sweetened beverages (per day), on average, in the last 7 days		75.3	84.4	82.9	81.9
Drinking energy drinks or shots on 0 of the past 30 days		61.6	61.9	57.4	61.4
Drinking energy drinks or shots on 1-4 of the past 30 days		18.1	20.0	23.0	19.9
Drinking energy drinks or shots on 5 or more of the past 30 days		20.3	18.1	19.7	18.7
Eating 3 or more servings of fruits (per day), on average, in the last 7 days		18.1	14.5	13.8	15.2
Eating 3 or more servings of vegetables (per day), on average, in the last 7 days		21.8	17.8	21.0	18.9

Estimated Body Mass Index (BMI): This figure incorporates reported height and weight to form a general indicator of physical health. Categories defined by The World Health Organization (WHO) 2000, reprinted 2004. Obesity: Preventing and Managing the Global Epidemic. WHO Tech Report Series: 894.

				Trans/	
	Ci	s Men	Cis Women	Gender Non-	Total
BMI	Percent (%)			conforming	
<18.5 Underweight		3.4	4.3	3.4	4.0
18.5-24.9 Healthy Weight		45.5	49.5	42.0	47.7
25-29.9 Overweight		25.1	21.6	25.2	22.7
30-34.9 Class I Obesity		12.3	14.3	17.6	14.4
35-39.9 Class II Obesity		8.5	5.5	6.7	6.3
≥40 Class III Obesity		5.1	4.9	5.0	5.0
Mean		26.9	26.4	27.3	26.6
Median		25.1	24.5	25.8	24.8
Std Dev		7.0	6.8	7.4	6.9

Students meeting the recommended guidelines for physical activity

Based on: US Dept of Health and Human Services. *Physical Activities Guidelines for Americans*, 2nd edition. Washington, DC: US Dept of Health and Human Services; 2018

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

Definitions:

- Recommendation for aerobic activity: 150 minutes or more of moderate-intensity physical activity per week or 75 minutes of vigorous-intensity
 physical activity or the equivalent combination
- Recommendation for strength training: 2 or more days a week of moderate or greater intensity activities that involve all major muscle groups
- Active Adults meet the recommendation for strength training AND aerobic activity
- Highly Active Adults meet the recommendation for strength training and TWICE the recommendation for aerobic activity (300 minutes or more of moderate-intensity physical activity per week or 150 minutes of vigorous-intensity physical activity or the equivalent combination)

			Trans/	
	Cis M Percent (%)	en Cis Women	Gender Non- contorming	Total
Guidelines met for aerobic exercise only	73.	9 64.0	66.4	66.5
Guidelines met for Active Adults	47.	9 33.2	17.2	34.7
Guidelines met for Highly Active Adults	36.	1 23.7	9.8	24.9

Food Security

Based on responses to the US Household Food Security Survey Module: Six-Item Short Form (2012) from the USDA Economic Research Service.

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
High or marginal food security (score 0-1)		49.0	47.5	36.9	46.4
Low food security (score 2-4)		29.3	27.4	28.7	27.9
Very low food security (score 5-6)		21.8	25.1	34.4	25.7
Any food insecurity (low or very low food security)		51.0	52.5	63.1	53.6

C. Health Care Utilization

				Trans/	
College students reported:		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Receiving psychological or mental health services within the last 12 months		27.6	34.1	49.2	34.3
Visiting a medical provider within the last 12 months		64.4	78.9	83.1	76.1

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Ever prescribed medication for a mental health condition		28.9	43.5	66.9	43.0
Prescribed before starting at current college/university*		23.8	34.5	60.5	35.0
Prescribed after starting at current college/university*		4.6	9.0	6.5	7.7
Last 12 months, prescribed medication for a mental health condition*		13.8	23.7	33.9	22.5
*These figures use all students in the sample as the denominator, rather than just those students					

who have ever been prescribed medication for a mental health condition			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
Percent (%)			conforming	
Ever had counseling for a mental health condition	34.3	47.0	71.5	46.8
Started counseling before starting at current college/university*	28.5	39.5	64.5	39.7
Started counseling after starting at current college/university*	5.9	7.4	6.5	6.9
Last 12 month, had counseling for a mental health condition*	14.6	26.0	37.1	24.6

^{*}These figures use all students in the sample as the denominator, rather than just those students who have ever had counseling for a mental health condition

College students reported:				Trans/	
	Percent (%)	Cis Men	Cis Women	Gender Non- conforming	Total
Flu vaccine within the last 12 month		38.7	39.0	39.5	38.9
Not starting the HPV vaccine series		13.0	13.2	10.5	13.1
Starting, but not completing HPV vaccine series		3.3	7.0	5.6	6.0
Completing HPV vaccine series		46.0	44.7	43.5	44.8
Not knowing their HPV vaccine status		37.7	35.1	40.3	36.1
Ever having a GYN visit or exam (females only)			39.8	36.3	
Having a dental exam in the last 12 months		76.1	77.1	73.4	76.4
Being tested for HIV within the last 12 months		7.1	8.1	11.3	8.2
Being tested for HIV more than 12 months ago		5.5	2.7	6.5	3.7
Wearing sunscreen usually or always when outdoors		15.1	32.5	26.6	28.1
Spending time outdoors with the intention of tanning at least once in the last 12 months		46.6	65.3	29.5	57.2

D. Impediments to Academic Performance

Respondents are asked in numerous places throughout the survey about issues that might have negatively impacted their academic performance within the last 12 months. This is defined as negatively impacting their performance in a class or delaying progress towards their degree. Both types of negative impacts are represented in the figures below. Please refer to the corresponding Data Report for specific figures on each type of impact. Figures in the left columns use all students in the sample as the denominator. Figures in the right columns use only the students that experienced that issue (e.g. students who used cannabis, reported a problem or challenge with finances, or experienced a particular health issue) in the denominator. (items are listed in the order in which they appear in the survey)

Negatively impacted academic performance	:e
among <u>all students in the sample</u>	

	Cis Men	Cis Women	Trans/ Gender Non-	Total
Percen		CIS VVOINCI	conforming	Total
Alcohol use	2.1	1.1	2.4	1.4
Cannabis/marijuana use	3.3	1.6	2.4	2.1
Pregnancy or a partner's pregnancy	0.0	0.4	0.0	0.3
Problems or challenges in the last 12 months				
Career	8.4	8.2	12.9	8.9
Finances	13.8	15.0	22.6	15.5
Procrastination	50.2	49.1	69.4	51.3
Faculty	6.3	5.1	8.9	5.8
Family	7.9	11.6	14.5	11.1
Intimate Relationships	10.0	9.2	8.9	9.3
Roommate/housemate	5.4	5.8	4.8	5.6
Peers	4.6	4.4	5.6	4.7
Personal appearance	4.6	6.9	12.9	7.1
Health of someone close to me	7.9	9.0	13.7	9.3
Death of a family member, friend, or someone clos	e			
to me	5.9	10.8	11.3	9.8
Bullying	2.5	2.0	4.0	2.4
Cyberbullying	0.8	0.9	1.6	1.1
Hazing	0.4	0.3	0.0	0.4
Microaggression	2.1	2.0	4.0	2.4
Sexual Harassment	1.7	2.8	4.0	2.8
Discrimination	1.7	1.3	3.2	1.7

Negatively impacted academic performance among only students that experienced the issue

Cis Men	Cis Women	Trans/ Gender Non-	Total
CIS IVICII	CIS WOITIEII	conforming	TOtal
3.5	1.7	3.5	2.3
11.1	6.0	5.4	7.0
11.1	0.0	5.1	,.0
0.0	42.9	0.0	42.9
30.3	29.8	32.0	30.6
28.0	28.0	35.4	28.9
62.8	63.2	76.1	64.9
57.7	52.8	50.0	53.3
31.1	28.1	24.3	27.9
27.0	26.3	26.2	26.5
22.8	16.4	20.7	17.9
23.4	15.7	14.3	16.8
8.7	10.9	17.8	11.4
20.7	20.6	24.3	21.2
26.4	39.2	48.3	37.8
31.6	25.9	23.8	27.3
28.6	20.0	25.0	23.5
100.0	100.0	0.0	66.7
13.2	15.3	9.1	14.0
33.3	24.1	21.7	25.2
19.0	20.4	14.3	19.2

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

Negatively impacted academic performance among <u>all students in the sample</u>

			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
Percent (%)			conforming	
Acute Diagnoses in the last 12 months				
Bronchitis	1.3	2.2	0.8	1.8
Chlamydia	8.0	0.0	0.0	0.2
Chicken Pox (Varicella)	0.0	0.1	0.0	0.1
Cold/Virus or other respiratory illness	12.6	19.8	29.0	19.2
Concussion	0.8	2.6	0.8	2.0
Gonorrhea	0.0	0.0	0.0	0.0
Flu (influenza or flu-like illness)	4.2	6.9	8.9	6.4
Mumps	0.0	0.0	0.0	0.0
Mononucleosis (mono)	0.4	1.2	1.6	1.1
Orthopedic injury	2.5	3.1	3.2	3.0
Pelvic Inflammatory Disease	0.0	0.1	0.0	0.1
Pneumonia	8.0	1.3	1.6	1.3
Shingles	0.0	0.1	0.0	0.1
Stomach or GI virus or bug, food poisoning or gastritis	4.2	6.6	3.2	5.6
Urinary tract infection	0.0	4.6	2.4	3.3

	G1 1.4	5 1 111	Trans/	
	Cis Men	Cis Women	Gender Non-	lotal
Percent (%)			conforming	
Any ongoing or chronic medical conditions diagnosed				
or treated in the last 12 months	22.2	35.0	61.3	34.9
Other impediments to academic performance				
Assault (physical)	0.8	1.3	0.8	1.2
Assault (sexual)	0.4	3.1	5.6	2.8
Allergies	5.0	4.7	4.8	4.7
Anxiety	26.8	39.5	62.1	38.9
ADHD or ADD	12.6	15.6	36.3	17.1
Concussion or TBI	0.8	3.0	0.8	2.2
Depression	21.3	27.0	58.1	29.0
Eating disorder/problem	8.0	3.2	7.3	3.1
Headaches/migraines	4.2	16.3	24.2	14.4
Influenza or influenza-like illness (the flu)	3.8	5.4	5.6	5.0
Injury	2.5	2.6	4.0	2.7
PMS	0.0	15.8	17.7	12.4
PTSD	0.4	3.9	8.1	3.6
Short-term illness	3.3	6.5	8.9	6.1
Upper respiratory illness	5.4	10.0	15.3	9.5
Sleep difficulties	16.7	26.8	50.0	26.9
Stress	34.7	42.6	64.5	43.1

Negatively impacted academic performance among only students that experienced the issue

		Trans/	
Cis Men	Cis Women	Gender Non-	Total
		conforming	
50.0	50.0	14.3	44.4
50.0	0.0	0.0	16.7
0.0	50.0	0.0	25.0
42.9	45.4	62.1	47.1
33.3	55.9	100.0	53.7
0.0	0.0	0.0	0.0
37.0	54.3	73.3	52.9
0.0	0.0	0.0	0.0
25.0	60.0	66.7	54.5
28.6	30.7	30.8	30.3
0.0	50.0	0.0	33.3
66.7	62.5	50.0	60.9
0.0	50.0	0.0	33.3
62.5	59.0	25.0	54.8
0.0	33.0	20.0	30.6

Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
31.5	43.2	67.9	44.0
25.0	43.5	20.0	36.1
12.5	37.7	63.6	38.8
13.5	10.8	9.0	11.0
44.1	49.2	69.4	50.9
58.8	66.7	77.6	67.3
22.2	64.7	25.0	53.2
58.6	60.1	69.2	61.6
11.1	16.1	25.7	17.3
16.1	35.7	41.1	33.8
24.3	38.5	36.8	35.0
17.1	24.4	31.3	23.3
0.0	36.9	38.6	36.9
14.3	42.6	38.5	38.8
22.9	33.8	34.4	32.4
23.2	30.2	32.2	29.4
38.5	46.9	63.9	47.9
48.8	51.5	70.8	53.4

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

E. Violence, Abusive Relationships, and Personal Safety

Within the last 12 months, college students reported experiencing:			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
Percent (%)			conforming	
A physical fight	3.3	1.1	0.0	1.5
A physical assault (not sexual assault)	1.3	2.3	3.3	2.3
A verbal threat	17.2	11.2	17.2	13.3
Sexual touching without their consent	2.9	5.0	5.7	4.6
Sexual penetration attempt without their consent	0.0	2.6	3.3	2.1
Sexual penetration without their consent	0.4	2.3	1.6	1.8
Being a victim of stalking	1.3	5.8	8.1	5.1
A partner called me names, insulted me, or put me down to make me feel bad	10.1	10.2	10.7	10.2
A partner often insisted on knowing who I was with and where I was or tried to limit my contact				
with family or friends	8.0	7.3	0.8	6.8
A partner pushed, grabbed, shoved, slapped, kicked, bit, choked or hit me without my consent	1.7	4.3	2.5	3.7
A partner forced me into unwanted sexual contact by holding me down or hurting me in some way	0.4	3.0	0.0	2.3
A partner pressured me into unwanted sexual contact by threatening me, coercing me, or using				
alcohol or other drugs	3.0	5.3	1.6	4.5

College students reported feeling **very safe***:

			Trans/	
Percent (%)	Cis Men	Cis Women	Gender Non- conforming	Total
On their campus (daytime)	87.0	85.8	86.3	85.8
On their campus (nighttime)	36.8	11.1	16.1	17.1
In the community surrounding their campus (daytime)	36.2	25.0	23.3	27.4
In the community surrounding their campus (nighttime)	12.2	5.2	1.8	6.3

 $[\]ensuremath{^{*}}\mbox{Note:}$ "Does not apply" responses were removed from the total in the denominator

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

F. Tobacco, Alcohol, and Other Drug Use

Ever Used

*Used in the last 3 months

		Evel C	ocu .	
	Trans/			
Percent (%)	Cis Men	Cis Women	Gender Non- conforming	Total
Tobacco or nicotine delivery products (cigarettes, e-				
cigarettes, Juul or other vape products, water pipe or				
hookah, chewing tobacco, cigars, etc.)	28.5	27.9	30.6	28.2
Alcoholic beverages (beer, wine, liquor, etc.) Cannabis (marijuana, weed, hash, edibles, vaped	59.4	62.4	65.0	61.9
cannabis, etc.) [Please report nonmedical use only.]	37.2	32.4	54.8	35.9
Cocaine (coke, crack, etc.)	1.7	1.4	2.4	1.6
Prescription stimulants (Ritalin, Concerta, Dexedrine,				
Adderall, diet pills, etc.) [Please report nonmedical use				
only.]	5.4	3.2	4.8	4.0
Methamphetamine (speed, crystal meth, ice, etc.)	0.0	0.4	1.6	0.5
Inhalants (poppers, nitrous, glue, gas, paint thinner,				
etc.)	4.2	1.8	4.1	2.5
Sedatives or Sleeping Pills (Valium, Ativan, Xanax, Klonopin, Librium, Rohypnol, GHB, etc.) [Please report				
nonmedical use only.]	2.5	3.2	4.0	3.2
Hallucinogens (Ecstasy, MDMA, Molly, LSD, acid,				
mushrooms, PCP, Special K, etc.)	7.9	4.6	9.8	5.9
Heroin	0.4	0.0	0.8	0.2
Prescription opioids (morphine, codeine, fentanyl,				
oxycodone [OxyContin, Percocet], hydrocodone				
[Vicodin], methadone, buprenorphine [Suboxone],				
etc.) [Please report nonmedical use only.]	3.4	1.9	1.6	2.3

		Trans/	
Cis Men	Cis Women	Gender Non- conforming	Total
19.2	19.7	16.9	19.2
51.0	56.5	54.8	54.7
27.6	23.2	41.1	26.0
0.0	0.1	0.0	0.1
2.1	2.0	1.6	2.1
0.0	0.4	0.0	0.4
1.7	0.7	1.6	1.0
0.0	1.8	0.0	1.3
2.1	1.5	2.4	1.8
0.0	0.0	0.0	0.0
0.8	0.7	0.8	0.8

Substance Specific Involvement Scores (SSIS) from the ASSIST

*Moderate risk use of the substance

*High risk use of the substance

Trans/

	Percent (%)	Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
Tobacco or nicotine delivery products	r crecine (70)	15.1	12.9	13.7	13.4
Alcoholic beverages		9.6	8.1	9.7	8.6
Cannabis (nonmedical use)		15.5	14.8	29.0	16.5
Cocaine		0.0	0.1	0.0	0.1
Prescription stimulants (nonmedical use)		8.0	0.9	2.4	1.1
Methamphetamine		0.0	0.3	1.6	0.4
Inhalants		1.3	0.4	1.6	0.7
Sedatives or Sleeping Pills (nonmedical use)		0.4	0.7	0.8	0.6
Hallucinogens		0.8	1.1	8.0	1.0
Heroin		0.0	0.0	8.0	0.1
Prescription opioids (nonmedical use)		0.8	0.4	0.8	0.6

Cis Men	Cis Women	Gender Non- conforming	Total
0.4	1.3	2.4	1.3
0.4	0.4	0.8	0.4
2.1	1.5	2.4	1.7
0.0	0.0	0.0	0.0
0.0	0.0	0.0	0.1
0.0	0.0	0.0	0.1
0.0	0.1	0.0	0.1
0.0	0.0	0.0	0.1
0.0	0.1	0.0	0.2
0.0	0.0	0.0	0.0
0.4	0.1	0.0	0.2

^{*}These figures use all students in the sample as the denominator, rather than just those students who reported lifetime use.

 $^{{}^*\!}T\!hese\ figures\ use\ all\ students\ in\ the\ sample\ as\ the\ denominator,\ rather\ than\ just\ those\ students\ who\ reported\ lifetime\ use.$

*Proportion of students who were prescribed a medication and used more than prescribed or more often than prescribed in the past 3 months

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Prescription stimulants		1.7	0.7	1.6	1.1
Prescription sedatives or sleeping pills		0.0	0.8	0.0	0.6
Prescription opioids		0.8	0.3	0.8	0.5

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

*Tobacco or nicotine delivery products used in the last 3 months

		Trans/		
	Cis Men	Cis Women	Gender Non-	Total
Percent (%)			conforming	
Cigarettes	9.6	5.7	12.1	7.2
E-cigarettes or other vape products (for example:				
Juul, etc.)	15.9	17.3	12.9	16.3
Water pipe or hookah	0.4	0.7	0.8	0.6
Chewing or smokeless tobacco	5.0	0.7	0.8	1.6
Cigars or little cigars	6.3	1.1	1.6	2.2
Other	0.8	0.1	0.0	0.3

^{*}These figures use all students in the sample as the denominator, rather than just those students who reported tobacco or nicotine delivery product use in the last 3 months.

Students in Recovery

2.4 % of college students surveyed (1.3 % cis men, indicated they were in recovery from alcohol or other drug use. 2.2 % cis women, and 5.6 % transgender/gender non-conforming)

When, if ever, was the last time you:

Drank Al	coho
----------	------

			Trans/	
Percent (%)	Cis Men	Cis Women	Gender Non- conforming	Total
Never	33.5	31.7	27.4	31.8
Within the last 2 weeks	37.7	36.9	33.9	36.6
More than 2 weeks ago but within the last 30 days	5.9	9.3	8.1	8.4
More than 30 days ago but within the last 3 months	11.3	9.8	12.9	10.4
More than 3 months ago but within the last 12	5.4	7.5	14.5	7.8
More than 12 months ago	6.3	4.7	3.2	4.9

^{*}Used Cannabis/Marijuana

		Trans/	
Cis Men	Cis Women	Gender Non- conforming	Total
58.6	66.1	41.9	61.8
16.7	13.0	25.0	15.1
3.3	3.9	4.8	3.9
3.3	5.8	11.3	5.9
8.8	5.0	6.5	6.0
9.2	6.2	10.5	7.3

Driving under the influence

• 10.0 % of college students reported driving after having any alcohol in the last 30 days.*

*Only students who reported driving in the last 30 days and drinking alcohol in the last 30 days were asked this question.

• 31.6 % of college students reported driving within 6 hours of using cannabis/marijuana in the last 30 days.*

*Only students who reported driving in the last 30 days and using cannabis in the last 30 days were asked this question.

Estimated Blood Alcohol Concentration (or eBAC) of college students. Due to the improbability of a student surviving a drinking episode resulting in an extremely high eBAC, all students with an eBAC of 0.50 or higher are also omitted from these eBAC figures. eBAC is an estimated figure based on the reported number of drinks consumed during the last time they drank alcohol in a social setting, their approximate time of consumption, sex, weight, and the average rate of ethanol metabolism. Only students who reported drinking alcohol within the last 3 months answered these questions.

^{*}These figures use all students in the sample as the denominator, rather than just those students who reported having a prescription. Note that the title of this table was changed in Fall 2022, but the figures remain the same.

^{*}Students were instructed to include medical and non-medical use of cannabis.

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
Estimated BAC	Percent (%)			conforming	
< .08		83.1	80.3	87.9	81.8
< .10		89.2	85.9	93.9	87.5
Mean		0.04	0.05	0.03	0.04
Median		0.01	0.02	0.01	0.02
Std Dev		0.06	0.06	0.04	0.06

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

^{*}Reported number of drinks consumed the last time students drank alcohol in a social setting.

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
Number of drinks	Percent (%)			conforming	
4 or fewer		71.8	78.8	86.8	78.2
5		3.1	9.1	5.9	7.5
6		7.6	4.1	2.9	4.7
7 or more		17.6	7.9	4.4	9.6
Mean		4.0	3.0	2.2	3.1
Median		2.0	2.0	1.0	2.0
Std Dev		3.8	2.3	2.2	2.7

^{*}Only students who reported drinking alcohol in the last three months were asked this question.

Reported number of times college students consumed $\underline{\text{five or more drinks}}$ in a sitting within the last two weeks:

Among all students surveyed

		Trans/		
	Cis Men	Cis Women		Total
Percent (%)			conforming	
Did not drink alcohol in the last two weeks (includes				
non-drinkers)	62.3	63.1	66.1	63.4
None	16.3	17.7	23.4	18.0
1-2 times	15.5	14.2	8.9	13.8
3-5 times	5.0	4.6	0.8	4.2
6 or more times	0.8	0.5	0.8	0.6

^{*}Only students who reported drinking alcohol in the last two weeks were asked this question.

*Among those who reported drinking alcohol within the last two weeks

Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
43.3	47.8	69.0	49.1
41.1	38.3	26.2	37.6
13.3	12.4	2.4	11.5
2.2	1.5	2.4	1.7

^{*}College students who drank alcohol reported experiencing the following in the last 12 months when drinking alcohol:

			Trans/	
Percent (Cis Men	Cis Women	Gender Non- conforming	Total
Did something I later regretted	15.3	15.8	10.5	15.0
Blackout (forgot where I was or what I did for a large period of time and cannot remember, even				
when someone reminds me)	8.3	10.7	8.1	9.9
Brownout (forgot where I was or what I did for short periods of time, but can remember once				
someone reminds me)	18.8	21.4	12.8	19.8
Got in trouble with the police	0.0	0.4	0.0	0.3
Got in trouble with college/university authorities	1.4	0.4	3.5	1.0
Someone had sex with me without my consent	0.7	1.5	2.4	1.4
Had sex with someone without their consent	0.0	0.4	0.0	0.3
Had unprotected sex	7.6	13.1	10.5	11.7
Physically injured myself	7.7	7.2	3.5	6.9
Physically injured another person	0.0	0.0	0.0	0.0
Seriously considered suicide	4.2	1.3	5.9	2.4
Needed medical help	0.0	1.1	0.0	0.7
Reported two or more of the above	18.6	22.2	17.1	20.8

^{*}Only students who reported drinking alcohol in the last 12 months were asked these questions.

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

G. Sexual Behavior

When, if ever, was the last time you had:

when, it ever, was the last time you had.		Olulis	CA	
Percent (%)	Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
Never	39.7	43.1	42.7	42.5
Within the last 2 weeks	26.4	28.6	28.2	28.1
More than 2 weeks ago but within the last 30 days	7.5	8.1	5.6	7.7
More than 30 days ago but within the last 3 months	7.5	6.9	7.3	7.0
More than 3 months ago but within the last 12	6.7	7.2	8.1	7.1
More than 12 months ago	12.1	6.1	8.1	7.7

Oral sex

Vaginal intercourse

Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
50.6	47.8	61.0	50.1
21.3	30.5	24.4	27.8
6.7	5.5	1.6	5.3
5.0	6.6	4.9	6.0
5.9	5.0	2.4	4.9
10.5	4.5	5.7	5.9

Anal intercourse

Percent (%)	Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
Never	78.1	88.0	82.8	85.2
Within the last 2 weeks	3.8	1.9	4.1	2.7
More than 2 weeks ago but within the last 30 days	2.1	0.7	1.6	1.1
More than 30 days ago but within the last 3 months	2.5	1.6	2.5	1.9
More than 3 months ago but within the last 12	5.1	2.7	3.3	3.3
More than 12 months ago	8.4	5.1	5.7	5.9

*College students who reported having oral sex, or vaginal or anal intercourse within the last 12 months reported having the following number of sexual partners:

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
None		0.0	0.7	0.0	0.5
1		63.1	73.8	85.5	72.7
2		18.0	12.1	8.1	12.9
3		7.4	7.0	3.2	6.9
4 or more		11.5	6.3	3.2	7.0
Mean		2.1	1.6	1.4	1.7
Median		1.0	1.0	1.0	1.0
Std Dev		2.8	1.9	1.2	2.1

^{*}Only students who reported having oral sex, or vaginal or anal intercourse in the last 12 months were asked this question.

College students who reported having oral sex, or vaginal or anal intercourse within the last 30 days who reported using a condom or another protective barrier most of the time or always:

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Oral sex		9.9	7.7	2.4	7.6
Vaginal intercourse		56.1	50.0	35.5	50.1
Anal intercourse		28.6	38.9	42.9	37.5

^{*}Only students who reported having oral sex, or vaginal or anal intercourse in the last 30 days were asked these questions.

College students who reported having vaginal intercourse (penis in vagina) within the last 12 months were asked if they or their partner used any method to prevent pregnancy the last time they had vaginal intercourse:

	Percent (%)	Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
Yes, used a method of contraception		79.6	84.4	70.7	82.4
No, did not want to prevent pregnancy		0.0	0.8	0.0	0.6
No, did not use any method		18.3	14.4	26.8	16.2
Don't know		2.2	0.3	2.4	0.8

^{*}Only students who reported having oral sex, or vaginal or anal intercourse in the last 12 months were asked this question.

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

*Those students who reported using a contraceptive use the last time they had vaginal intercourse, reported they (or their partner) used the following methods:

en Cis Women	Gender Non- conforming	Total
43.0	conforming	
43.0		
	31.0	41.2
5.0	10.3	5.5
5.4	6.9	6.5
1.3	3.4	1.5
1.7	6.9	1.7
7.0	3.4	6.9
3 10.7	20.7	10.7
61.4	48.3	62.5
0.7	0.0	0.5
0.3	0.0	0.2
0.0	0.0	0.0
32.2	24.1	29.8
11.4	20.7	10.9
0.3	6.9	1.0
0.7	0.0	0.7
45.0	48.3	45.4
00.0	2/1	27.5
	0.0 32.2 11.4 0.3 0.7	0.0 0.0 32.2 24.1 11.4 20.7 0.3 6.9 0.7 0.0

^{*}Only students who reported they or their partner used a method the last time they had vaginal intercourse were asked these questions. This question was select all that apply.

College students who reported having vaginal intercourse (penis in vagina) within the last 12 months were asked if they or their partner used emergency contraception ("morning after pill" or "Plan B") in the last 12 months:

Yes (17.2 % cis men, 24.4 % cis women, 22.0 % trans/gender non-conforming)

College students who reported having vaginal intercourse (penis in vagina) within the last 12 months were asked if they experienced an unintentional pregnancy or got someone pregnant within the last 12 months:

Yes (0 % cis men, 1.4 % cis women, 0.0 % trans/gender non-conforming)

H. Mental Health and Wellbeing

Kessler 6 (K6) Non-Specific Psychological Distress Score (Range is 0-24)

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
No or low psychological distress (0-12)		82.4	75.9	51.6	74.5
Serious psychological distress (13-24)		17.6	24.1	48.4	25.5
Mean		7.67	9.23	12.27	9.25
Median		7.00	9.00	12.00	9.00
Std Dev		4.87	4.97	4.66	5.06

UCLA Loneliness Scale (ULS3) Score (Range is 3-9)

			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
Percei	nt (%)		conforming	
Negative for Ioneliness (3-5)	43.9	47.1	31.5	44.6
Positive for Ioneliness (6-9)	56.1	52.9	68.5	55.4
Mean	5.68	5.65	6.32	5.74
Median	6.00	6.00	6.00	6.00
Std Dev	1.88	1.86	1.80	1.87

Diener Flourishing Scale - Psychological Well-Being (PWB) Score (Range is 8-56)

(higher scores reflect a higher level of psychological well-being)

. ,	U	O,			
				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
				conforming	
		44.10	44.95	40.09	44.17
		46.00	47.00	41.00	46.00
		8.44	7.81	8.75	8.20

Cis Men n = 239
Cis Women n = 742
Trans/GNC n = 124

The Connor-Davison Resilience Scale (CD-RISC2) Score (Range is 0-8)

(higher scores reflect greater resilience)

			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
			conforming	
Mean	6.03	5.80	5.23	5.78
Median	6.00	6.00	5.00	6.00
Std Dev	1.54	1.50	1.68	1.54

Self injury

Mean Median Std Dev

• 16.2 % of college students surveyed (10.9 % cis men, 16.0 % cis women, and 28.2 % trans/gender non-conforming) indicated they had intentionally cut, burned, bruised, or otherwise injured themselves within the last 12 months.

Within the last 12 months, have you had problems or challenges with any of the following:

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Academics		46.0	46.5	66.1	48.6
Career		27.7	27.7	40.3	29.2
Finances		49.8	53.9	63.7	54.2
Procrastination		80.8	78.0	91.1	79.9
Faculty		10.9	9.7	17.7	11.0
Family		25.6	42.0	59.7	40.5
Intimate relationships		37.2	35.1	33.9	35.5
Roommate/housemate		24.5	35.6	23.4	31.9
Peers		20.2	28.7	39.5	28.2
Personal appearance		53.6	63.7	72.6	62.7
Health of someone close to me		38.5	44.1	56.9	44.3
Death of a family member, friend, or some	one close				
to me		22.2	27.6	23.4	26.0
I was bullied		7.9	7.8	16.9	8.9
I was cyberbullied		2.9	4.7	6.5	4.6
I was hazed		0.4	0.3	1.6	0.5
I experienced microagression(s)		15.9	13.4	44.4	17.6
I was sexually harassed		5.0	11.9	18.5	11.3
I experienced discrimination		8.8	6.6	22.8	8.9
*Only students who reported a problem or challenge in	the last 12 mon	ths were asked	d about level of dis	stress.	
Students reporting none of the above		8.8	2.7	0.8	3.8
Students reporting only one of the above		7.9	6.5	4.0	6.7
Students reporting 2 of the above		10.9	10.5	2.4	9.7
Students reporting 3 or more of the above		72.4	80.3	92.7	79.9

*Of those reporting this issue, it caused moderate or high distress

		Trans/	
Cis Men	Cis Women	Gender Non- conforming	Total
76.4	82.7	92.7	83.1
59.1	71.4	70.0	68.8
68.9	76.9	81.0	75.8
59.9	65.2	76.1	65.3
41.7	45.1	45.5	44.5
60.7	65.3	65.8	64.7
61.8	61.2	73.8	62.8
51.7	56.9	48.3	55.7
33.3	48.1	53.1	47.1
48.4	61.1	61.1	58.5
52.2	59.0	68.6	59.3
75.5	67.0	82.8	70.4
57.9	58.6	52.4	57.6
57.1	44.1	37.5	46.0
0.0	100.0	0.0	50.0
31.6	37.1	43.6	38.0
50.0	52.3	69.6	55.7
47.6	49.0	67.9	54.5

Suicide Behavior Questionnaire-Revised (SBQR) Screening Score (Range is 3-18)

	Percent (%)	Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
Negative suicidal screening (3-6)	r creent (70)	66.8	68.2	38.2	64.4
Positive suicidal screening (7-18)		33.2	31.8	61.8	35.6
Mean		5.71	5.64	8.54	5.99
Median		4.00	4.00	8.00	5.00
Std Dev		3.40	3.37	3.86	3.56

Cis Men n = 239 Cis Women n = 742 Trans/GNC n = 124

Suicide attempt

• 2.9 % of college students surveyed

2.9 % cis men,

2.6 % cis women, and

• 4.8 % trans/gender non-conforming indicated they had attempted suicide within the last 12 months.

Within the last 30 days, how would you rate the overall level of stress experienced:

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
No stress		2.1	0.9	0.0	1.2
Low		27.2	16.1	10.5	17.8
Moderate		52.3	52.1	38.7	50.5
High		18.4	30.9	50.8	30.5

I. Acute Conditions

College students reported being diagnosed by a healthcare professional within the last 12 months with:

			Trans/	
Percent (%)	Cis Men	Cis Women	Gender Non- conforming	Total
Bronchitis	2.5	4.6	5.6	4.2
Chlamydia	1.7	1.1	0.8	1.2
Chicken Pox (Varicella)	0.9	0.3	0.0	0.4
Cold/virus or other respiratory illness (for example:				
sinus infection, ear infection, strep throat, tonsillitis,				
pharyngitis, or laryngitis)	29.3	44.0	46.8	41.1
Concussion	3.0	4.9	0.8	4.0
Gonorrhea	0.4	0.1	0.0	0.2
Flu (influenza) or flu-like illness	11.3	12.9	12.1	12.4
Mumps	8.0	0.3	0.0	0.4
Mononucleosis (mono)	2.1	2.1	2.4	2.1
Orthopedic injury (for example: broken bone,				
fracture, sprain, bursitis, tendinitis, or ligament injury)	8.8	10.1	10.5	9.8
Pelvic Inflammatory Disease	0.4	0.3	0.0	0.3
Pneumonia	1.3	2.2	3.2	2.1
Shingles	0.4	0.3	0.0	0.3
Stomach or GI virus or bug, food poisoning or gastritis	6.7	11.6	12.9	10.6
Urinary tract infection	0.8	14.5	12.4	11.3

Cis Men n = 239 Cis Women n = 742 Trans/GNC n = 124

J. Ongoing or Chronic Conditions

Mental Health

The questions for the *ongoing or chronic conditions* are presented differently in this report than the order they appear in the survey. In the survey, all items appear in a single list, ordered alphabetically. In this report, the conditions are presented in groups to ease burden on the reader. The findings are divided into mental health conditions, STIs and other chronic infections, and other ongoing or chronic conditions in this report.

College students reported <u>ever</u> being diagnosed with the following:

*Of those ever diagnosed, those reporting contact with healthcare or MH professional within last 12 months

Mental Health	uic	gnosca With	the following.	
			Trans/	
	Cis Men	Cis Women	Gender Non-	Total
Percent (%)			conforming	
ADD/ADHD - Attention Deficit/Hyperactivity				
Disorder	19.7	15.7	36.3	18.9
Alcohol or Other Substance Use Disorder	0.4	0.3	1.6	0.5
Anxiety (for example: Generalized Anxiety, Social				
Anxiety, Panic Disorder, Specific Phobia)	25.5	48.9	65.0	45.7
Autism Spectrum	7.1	3.5	16.1	5.8
Bipolar and Related Conditions (for example: Bipolar I,				
II, Hypomanic Episode)	2.5	3.8	5.6	3.7
Borderline Personality Disorder (BPD), Avoidant				
Personality, Dependent Personality, or another				
personality disorder	0.4	2.2	5.6	2.2
Depression (for example: Major depression, persistent				
depressive disorder, disruptive mood disorder)	22.6	35.8	60.2	35.6
Eating Disorders (for example: Anorexia Nervosa,				
Bulimia Nervosa, Binge-Eating)	1.3	8.5	13.7	7.6
Gambling Disorder	0.4	0.0	0.0	0.1
Insomnia	3.0	8.6	22.8	9.0
Obsessive-Compulsive and Related Conditions (for				
example: OCD, Body Dysmorphia, Hoarding,				
Trichotillomania and other body-focused repetitive				
behavior disorders)	3.3	8.8	14.5	8.3
PTSD (Posttraumatic Stress Disorder), Acute Stress				
Disorder, Adjustment Disorder, or another trauma- or				
stressor- related condition	2.9	8.2	18.5	8.3
Schizophrenia and Other Psychotic Conditions (for				
example: Schizophrenia, Schizoaffective Disorder,				
Schizophreniform Disorder, Delusional Disorder)	0.0	0.0	2.4	0.3
Tourette's or other neurodevelopmental condition				
not already listed	0.8	0.5	3.2	1.0
Traumatic brain injury (TBI)	1.3	0.7	0.8	0.8
, , , ,				

^{*}Only students who reported ever being diagnosed were asked about contact with a healthcare or mental health professional within the last 12 months.

Percent (%)	Cis Men	Cis Women	Trans/ Gender Non- conforming	Total
Students reporting none of the above	57.3	43.5	21.8	44.0
Students reporting only one of the above	15.9	16.8	13.7	16.3
Students reporting both Depression and Anxiety	17.6	32.8	54.0	31.9
Students reporting any two or more of the above				
(excluding the combination of Depression and Anxiety)	9.2	6.9	10.5	7.8

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

College students reported ever being diagnosed with the following:

*Of those ever diagnosed, had contact with healthcare or MH professional within last 12 months

STI's/Other chronic infections		diagnosed with the following:			
		Trans/		Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Genital herpes		0.4	0.1	1.6	0.4
Hepatitis B or C		0.4	0.1	0.0	0.2
HIV or AIDS		0.4	0.0	0.0	0.1
Human papillomavirus (HPV) or genital warts	5	0.4	0.7	0.0	0.5

		Trans/	
Cis Men	Cis Women	Gender Non-	Total
		conforming	
100.0	100.0	100.0	100.0
100.0	0.0	0.0	50.0
100.0	0.0	0.0	100.0
100.0	80.0	0.0	83.3

^{*}Only students who reported ever being diagnosed were asked about contact with a healthcare or mental health professional within the last 12 months.

	College students reported ever being
her Chronic /Ongoing Medical Conditions	diagnosed with the following:

*Of those ever diagnosed, had contact with healthcare or MH professional within last 12 months

Other Chronic /Ongoing Medical Conditions	dis	agnosed with	the following:			mo	nth	nths
Galer Chrome / Origonia Medical Conditions	uia	61103CU WILLI	Trans/	_		mont		Trans/
Percent (%)		Cis Women	Gender Non- conforming	Total	Cis Men	Cis Women		
ne	24.7	29.5	31.5	28.7	32.8	34.0		26.3
rgies - food allergy	7.9	14.6	18.9	13.5	26.3	29.5		30.4
ies - animals/pets	15.5	12.8	19.4	14.1	21.6	33.3		20.8
ies - environmental (for example: pollen, grass,								
mold)	24.3	31.0	38.7	30.3	20.7	32.0		27.7
a	18.8	17.3	21.0	18.0	31.1	41.7		34.6
	0.4	0.8	0.0	0.6	100.0	66.7		0.0
lisease	0.0	1.5	1.6	1.2	0.0	63.6		100.0
pain (for example: back or joint pain, arthritis,								
pain)	2.9	10.7	21.1	10.1	71.4	73.1		65.4
or pre-diabetes/insulin resistance	1.3	4.1	6.5	3.7	33.3	90.0		62.5
triosis	0.0	2.7	3.2	2.2	0.0	70.0		50.0
sophageal Reflux Disease (GERD) or acid								
	2.1	7.7	14.5	7.2	40.0	57.9		50.0
vascular disorders (for example: atrial								
on or other cardiac arrhythmia, mitral valve								
se or other valvular heart disease, congenital								
ndition)	0.8	3.2	3.2	2.7	50.0	79.2		50.0
ood pressure (hypertension)	3.8	3.7	1.6	3.5	77.8	59.3		50.0
olesterol (hyperlipidemia)	3.8	2.6	0.0	2.5	44.4	63.2		0.0
bowel syndrome (spastic colon or spastic								
	2.5	7.9	12.9	7.3	83.3	57.9		25.0
e headaches	5.5	17.9	22.0	15.8	23.1	61.5		44.4
ic Ovarian Syndrome (PCOS)	0.0	6.9	12.1	6.0	0.0	80.4		73.3
onea	2.9	2.8	3.2	3.0	28.6	38.1		100.0
condition or disorder	0.0	4.2	5.7	3.4	0.0	87.1		57.1
system disorder (for example: bladder or								
disease, urinary reflux, interstitial cystitis)	0.0	2.7	4.0	2.3	0.0	45.0		60.0

^{*}Only students who reported ever being diagnosed were asked about contact with a healthcare or mental health professional within the last 12 months.

Cis Men n =	239
Cis Women n =	742
Trans/GNC n =	124

Students who reported being diagnosed with diabetes or pre-diabetes/insulin resistance, indicated they had:

				Trans/	
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Type I Diabetes		0.0	27.3	0.0	18.8
Type II Diabetes		0.0	17.4	0.0	12.1
Pre-diabetes or insulin resistance		66.7	72.4	100.0	77.5
Gestational Diabetes		0.0	4.5	0.0	3.1

K. Sleep

Reported amount of time to usually fall asleep at night (sleep onset latency):

				Trans/	
	Percent (%)	Cis Men	Cis Women	Gender Non- conforming	Total
Less than 15 minutes		44.4	36.4	26.6	36.7
16 to 30 minutes		33.1	30.0	24.2	29.9
31 minutes or more		22.6	33.6	49.2	33.3

Over the last 2 weeks, students reported the following average amount of sleep (excluding naps):

		On weeknights			
		Trans/			
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
Less than 7 hours		36.8	40.6	46.8	40.6
7 to 9 hours		61.5	57.0	50.0	57.1
10 or more hours		1.7	2.4	3.2	2.3

Λ-	14100	lond	nights
OII	wee	Kenu	1112116

		Trans/	
Cis Men	Cis Women	Gender Non-	Total
		conforming	
21.3	20.5	15.3	20.2
67.8	67.2	64.5	66.9
10.9	12.3	20.2	12.8

Students reported the following on 3 or more of the last 7 days:

Ealt tired	or cloopy	during the	day
reit tirea	or sieeby	during the	aav

			Trans/		
		Cis Men	Cis Women	Gender Non-	Total
	Percent (%)			conforming	
0 days		5.9	1.9	0.0	2.6
1-2 days		31.8	13.1	8.9	16.6
3-5 days		47.3	44.9	31.5	44.0
6-7 days		15.1	40.1	59.7	36.8

Got enough sleep so that they felt rested

		Trans/	
Cis Men	Cis Women	Gender Non-	Total
		conforming	
11.3	18.8	27.6	18.2
38.5	43.9	43.9	42.7
39.7	29.3	22.8	30.8
10.5	8.0	5.7	8.3

Demographics and Sample Characteristics

		2		
• Age	73.2 %	Students describe themselv Straight/Hotorosovual:	/es as 60.5	0/
18 - 20 years:		Straight/Heterosexual:		
21 - 24 years:	24.6 %	Asexual:	4.0	
25 - 29 years:	1.3 %	Bisexual:	14.8	
30+ years:	1.0 %	Gay:	2.9	
Mean age: 19.8 years		Lesbian:	5.4	
Median age: 19.0 years		Pansexual:	5.9	
		Queer:	3.7	
• Gender*		Questioning:	2.2	
Cis Women:	66.4 %	Identity not listed above:	0.7	%
Cis Men:	21.4 %			
Fransgender/Gender Non-conforming:	11.1 %	• Housing	,	•
See note on page 2 regarding gender categories		Campus or university housing		
		Fraternity or sorority residen		
Student status		Parent/guardian/other family		
st year undergraduate:	40.0 %	Off-campus:	30.2	
2nd year undergraduate:	23.2 %	Temporary or "couch surfing"		
Brd year undergraduate:	19.7 %	Don't have a place to live:	0.1	
4th year undergraduate:	14.5 %	Other:	0.2	%
th year or more undergraduate:	2.3 %			
Master's (MA, MS, MFA, MBA, etc.):	0.0 %	 Students describe themselven 	es as**	
Doctorate (PhD, EdD, MD, JD, etc.):	0.0 %	American Indian or Native Al	askan: 2.0	%
Not seeking a degree:	0.0 %	Asian or Asian American:	4.4	%
Other:	0.3 %	Black or African American:	8.6	%
		Hispanic or Latino/a/x:	7.8	%
Full-time student:	98.4 %	Middle Eastern/North Africa	n (MENA)	
Part-time student:	1.0 %	or Arab Origin:	0.4	%
Other student:	0.6 %	Native Hawaiian or Other		
		Pacific Islander Native:	0.4	%
Relationship status		White:	82.5	%
Not in a relationship:	51.4 %	Biracial or Multiracial:	5.2	%
n a relationship, not married/partnered:	46.6 %	Identity not listed above:	0.6	%
Married/partnered:	2.0 %	,		
•		If Hispanic or Latino/a/x, are y	ou**	
Primary Source of Health Insurance**		Mexican, Mexican American,		%
College/university sponsored SHIP plan:	2.6 %	Puerto Rican:	11.5	%
Parent or guardian's plan:	77.9 %	Cuban:	1.1	%
Employer (mine or my spouse/partners):	0.6 %	Another Hispanic, Latino/a/x	, or	
Medicaid, Medicare, SCHIP, or VA:	12.4 %	Spanish Origin:	23.0	%
Bought a plan directly or from the exchange:	0.3 %	_		
Embassy/sponsoring agency (international):	0.0 %	If Asian or Asian American, are	you**	
Another source:	0.6 %	East Asian:	57.1	%
Have insurance, but don't know source:	3.0 %	Southeast Asian:	20.4	
Don't have health insurance:	4.3 %	South Asian:	16.3	
Don't know if I have health insurance:	2.3 %	Other Asian:	8.2	
Students that selected more than one source:	4.5 %	2 55. 7 556111	5.2	
stadents that selected more than one source.	7.5 /0	 Visa status & location of st 	udv	
Student Veteran:	1.6 %	- Tiba Status & location of St	Do not have/need a U.S.	Have/need
- Staucht Veteran.	1.0 /0		visa	U.S. visa
Parent or primary responsibility for someone		Studying		
else's child/children under 18 years old:	1.8 %	in U.S.	79.4 %	14.3 %
	2.2 /0	Studying		
**this question was select all that apply, totals may add up to over 100%		Outside U.S.	5.6 %	0.6 %
and question reas select an trial apply, totals may dud up to over 100%		3 d t 3 d 5 d 5 d 5 d 5 d 5 d 5 d 5 d 5 d 5 d		17

First generation students:	43.8 %	 Participated in organized college athletics 		
(Proportion of students for whom no		Varsity:	3.2 %	
parent/guardian have completed a		Club sports:	8.8 %	
bachelor's degree)		Intramurals:	6.7 %	
• Do you have any of the following?		• Member of a <u>social</u> fraternity or sorority		
Attention Deficit/Hyperactivity Disorder		Greek member:	6.5 %	
(ADD or ADHD):	22.6 %			
Autism Spectrum Disorder:	10.0 %	Dropout Intention		
Deaf/Hearing loss:	2.9 %	Very, moderately, or slightly likely to:		
Learning disability:	7.2 %	Leave school and transfer schools:	8.5 %	
Mobility/Dexterity disability:	4.2 %	Leave school without transferring:	4.6 %	
Blind/low vision:	4.0 %			
Speech or language disorder:	1.5 %			